

# PERK

PLASTIC SURGERY  
BEVERLY HILLS

## POST-OP | *Facial Implants*

(310) 400-6180

concierge@perkplasticsurgery.com

### **What will I see when I look into a mirror?**

Depending on the type of implant, you may have incisions underneath your chin, inside your mouth or under your lash line. For the incision inside the mouth (if you have one). Rinse the inside of the mouth after each meal or three times per day with ½ strength Listerine.

For incisions around your lash line: please apply the antibiotic ointment prescribed by the office to the lower lash line.

For incisions underneath your chin: please leave the dressing in place.

You will have bruising and swelling around your face. You will have a compression garment around your face. You will wear this compression garment until seen in the office.

You will feel numbness on your face. It will gradually improve over the next couple of weeks. You will feel a tingling/burning sensation of your face as the nerves to your face heal from surgery.

### **When can I shower or take a bath?**

You can shower 24 hours after surgery. Wipe your face. Don't let the water hit your face. Don't get the compression garment around your jaw wet.

### **What can I do after surgery?**

Ice the area 20 minutes on and 10 minutes off while you are awake for the first three days. The more you can take it easy and ice the area, the less swelling there will be.

Sleep on your back with your head elevated at approximately a 45 degree angle. 2-3 pillows

Do not wear clothing that pulls over the head.

Avoid bending over for long periods of time.

You should "take it easy" the day of surgery and gradually increase your activity as you feel more comfortable. Avoid heavy exercise also during this time and only gradually return to your normal routine. No strenuous activity or heavy lifting for 4 weeks. You can resume all of your normal activities about four weeks after surgery.

### **When can I drive?**

You should never drive if you are still taking any pain medication other than Tylenol.

# PERK

PLASTIC SURGERY  
BEVERLY HILLS

## **When will I be able to return to work?**

Depending on the type of work you do, you should be able to return to your job as soon as you are comfortable.

## **How much pain will I have?**

The most discomfort that you will have lasts 5-7 days, sometimes a little longer. As you feel more comfortable, your need for medication will be less.

## **What should I eat after surgery?**

A light diet is best for the day of surgery. Begin by taking liquids slowly and progress to soups or jello.

A soft diet that does not require hard chewing is encouraged for 1 week.

Avoid salty foods. Avoid spicy foods.

If there is an incision inside your mouth, be VERY CAREFUL when you brush your teeth. Brush the top teeth only and back bottom teeth.

## **“What medications should I use?”**

Pain medication: I will prescribe a pain medication for you. Follow the directions on the bottle for their use. The pain medication will make you feel drowsy. Have someone assist you in your home and do not attempt to drive while you are taking the pain medication.

- A few days after surgery most women will find that Tylenol will take care of the discomfort.
  - Do not use aspirin for five days after the surgery.
  - It is best to take pain medication with crackers, jello, etc.
  - If you have no pain, do not take the medication.
- Alcohol and pain medication should not be taken together.

Antibiotic: You will be given an antibiotic. Please follow the instructions from your pharmacist or that are written on the packaging.

Stool softener: Narcotic pain medications can sometimes cause constipation. An over-the-counter stool softener (Colace) is recommended if you are prone to this.

- Prune juice mixed with 7-Up (half and half) for mild constipation
- If severely constipated use Miralax first as it is a mild but effective laxative, Phillips Milk of Magnesia, and fleet enemas (regular or oil retention) as a last resort.

Anti-nausea medication: Some nausea is normal in the first 24-48 hours following surgery. If you are nauseated, please take the anti-nausea medication.

You may have additional medications.

# PERK

PLASTIC SURGERY  
BEVERLY HILLS

## “Can I drink alcohol?”

Do not drink alcohol while taking narcotic pain medication after your surgery.

## “When will I be seen in the office after surgery?”

You will normally be seen in the office within one week of your surgery.

## “Should I apply anything to the incisions to help make it scar better?”

We will give you instructions on how to improve the scar appearance.

## “How will I know if I am having a problem?”

I NEED TO KNOW ABOUT THESE PROBLEMS IMMEDIATELY:

These are the complications that will require a change in your post-operative care!

**Hematoma** (blood collecting under your incisions) can occur within a few days of surgery. The warning signs are:

- Significant swelling
- Excessive or growing bruising

**Infection** is rare. It is normal to have a very small amount of drainage from your incisions for one to 2 days.  
Signs of infection are:

- Increased temperature
- Increasing drainage from the incisions
- Increasing redness around the incisions

**Leg swelling** with or without associated pain may indicate a problem with the vein circulation in your leg(s). Although some swelling is to be expected due to your surgery, intravenous fluids given to you during the surgery and your decreased level of activity, this swelling is normally mild, painless and affects both legs evenly. If you have a large amount of leg swelling (either one or both legs) or if you experience pain in your legs contact the office immediately.

**Breathing problems** after surgery are rare but can be a serious complication. If you develop any chest and/or back pain or the feeling of being short of breath you must contact my office or be seen in the nearest emergency medical facility without delay.

**Medication reactions** may occur with the drugs prescribed for you. If you develop a skin rash, itching, vomiting, or diarrhea, stop taking your medication and contact my office.